


Standard Form to Vertex Form – Day 2

Quadratic Relations Concept Map



Recall When given a quadratic in the form $y = ax^2 + bx + c$ we complete the square to express it in the form $y = a(x - h)^2 + k$. This form is easier

to graph and analyze.

Example 1 a) Write the following equation in the form $y = a(x - h)^2 + k$.

$$y = x^2 - 8x + 19$$

$\nearrow \left(-\frac{-8}{2}\right)^2$
 $= (-4)^2$
 $= 16$

$$y = \underline{x^2 - 8x + 16} - 16 + 19$$

Add and subtract the square of half the co-efficient of x

Group the perfect square trinomial.

$$y = \underline{(x - 4)^2} - 16 + 19$$

Factor the perfect square trinomial (write as square of a binomial).

$$y = \underline{(x - 4)^2 + 3}$$

Simplify outside the brackets.

b) Analyze the relation.

opens:

up

stretch/comp:

none

vertex:

(4, 3)

eq'n of axis of sym.:

$x = 4$

values x may take:

$\mathbb{R}$ or $-\infty$ to $+\infty$


values y may take:

$\{y \geq 3, y \in \mathbb{R}\}$ or 3 to $+\infty$

max/min val.:

minimum, 3 "greater than or equal to"
 "is an element of"
 "the set of all real numbers"

c) Graph the relation:



Example 2 a) Write the following equation in the form $y = a(x - h)^2 + k$.

$$y = \underline{2x^2 + 4x - 1}$$

$$y = \underline{2(x^2 + 2x) - 1}$$
 Group the first two terms and factor out the co-efficient of x^2

$$y = \underline{2(x^2 + 2x + 1 - 1) - 1}$$
 Inside the brackets, add and subtract the square of half the co-efficient of x

$$y = \underline{2((x+1)^2 - 1) - 1}$$
 Factor the perfect square trinomial (first three terms inside brackets).


$$y = \underline{2(x+1)^2 - 2 - 1}$$
 Distribute the number outside the large brackets to terms inside large brackets.

$$y = \underline{2(x+1)^2 - 3}$$
 Simplify outside the brackets.

b) Analyze the relation.

opens: $\uparrow$
 stretch/comp: stretch, factor of 2
 vertex: $(-1, -3)$
 eq'n of axis of sym.: $x = -1$
 values x may take: $\mathbb{R}$ or
 values y may take: $\{y \geq -3, y \in \mathbb{R}\}$
 max/min val.: minimum, -3

c) Graph the relation:



Example 3 a) Write the following equation in the form $y = a(x - h)^2 + k$.

$$y = -3x^2 - 12x - 7$$

$$y = -3(x^2 + 4x) - 7$$

$$y = -3(\underline{x^2 + 4x + 4 - 4}) - 7$$

$$y = -3(\underline{(x+2)^2 - 4}) - 7$$

$$y = -3(x+2)^2 + 12 - 7$$

$$y = -3(x+2)^2 + 5$$

b) Analyze the relation.

opens: *down*
stretch/comp: *stretch, factor of 3*

vertex: *$(-2, 5)$*

eq'n of axis of sym.: *$x = -2$*


values x may take: $\mathbb{R}$

values y may take: *$\{y \leq 5, y \in \mathbb{R}\}$*

max/min val.:

maximum, 5

c) Graph the relation:



Opportunity to Learn Your OTL!

Complete all questions in the provided handout that accompanies this lesson.
Pg. 270-271 #7cde, 8, 12, 13 (use Nspire), 14