


# The Quadratic Formula

## Quadratic Relations Concept Map



## The Quadratic Formula

### Recall

We already know two ways to solve a quadratic equation.

1. Solve by graphing.

(look for where the graph crosses the  $x$ -axis)


$\therefore$  solutions are  $x = 2$  and  $x = 4$ .

2. Solve by factoring.

$$x^2 + 7x + 12 = 0$$

$$(x + 3)(x + 4) = 0$$

$\therefore$  solutions are  $x = -3$  and  $x = -4$ .



What happens when these methods don't work well?

For example, what are the solution(s) to the quadratic equation:

$$x^2 - 3x + 1 = 0$$

### The Quadratic Formula (when all else fails...)

Given a quadratic equation in the form:

$$ax^2 + bx + c = 0$$

... you can find the solutions / roots /  $x$ -intercepts / zeroes using this formula:

“Where does this formula come from?”

[You can find out here.](#)

**Example 1** Solve each quadratic equation.

a)  $4x^2 - 12x + 9 = 0$   
 $a =$

Remember...  $ax^2 + bx + c = 0$  so:  $b =$   
 $c =$

To solve, just sub into the quadratic formula:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

∴ the solution is

Consider: what does this mean? How many times does the graph of this quadratic cross the  $x$ -axis? (We'll come back to this later).

b)  $x^2 - 2x + 3 = 0$       So...       $a =$        $b =$        $c =$

Sub:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Consider: what does this mean? Where would the graph of this quadratic cross the  $x$ -axis?

c)  $x^2 - 3x + 1 = 0$  So...  $a =$   $b =$   $c =$

Sub:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$x = \quad \text{or} \quad x =$$

Sometimes we need approximate answers, so we keep going (rounding off the  $\sqrt{5}$ ).

$$\therefore \text{the solutions are } x = \quad \text{or} \quad x =$$

$$\text{or } x \doteq \quad \text{or } x \doteq$$


Consider: we have two solutions for  $x$ . What does this mean? How many times does this quadratic cross the  $x$ -axis?

**Follow-up: The Discriminant**


Look back at parts a), b), and c) in example 1. How many solutions were there in each case?

Part a) had one solution,

Graphically:




Part b) had no solutions. Graphically:



Part c) had two solutions,

or

Graphically:



We can predict the number of solutions by examining the discriminant.

What is the discriminant? It's the  $b^2 - 4ac$  part of the quadratic formula, in short, everything under the  $\sqrt{\phantom{x}}$  sign.

1. If  $b^2 - 4ac = 0$  there is one solution (technically, "two real but equal roots").

In part a), the discriminant is       , and sure enough, there was one solution.

2. If  $b^2 - 4ac < 0$  there are no solutions (technically, "no real roots").

In part b), the discriminant is       , and there were no solutions.

3. If  $b^2 - 4ac > 0$  there are two solutions (technically, "two real roots").

In part c), the discriminant is       , and there were two solutions.

That's it. Whew! All of this boils down to: when you can't solve a quadratic equation by graphing or by factoring, you can always break out the quadratic formula and solve it that way.

**Opportunity to Learn**

Complete all questions in the provided handout that accompanies this lesson.